



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Retention of solute and particle markers in the digestive tract of chinchillas (*Chinchilla laniger*)

Hagen, K B ; Dittmann, M T ; Ortmann, S ; Kreuzer, Michael ; Hatt, Jean-Michel ; Clauss, Marcus

Abstract: The chinchilla (*Chinchilla laniger*) is a herbivorous hystricomorph South American rodent for which no mean digesta retention times have been reported so far. Six animals (mean body mass \pm standard deviation: 513 ± 99 g) on a diet of grass hay and lucerne-based pellets were given a pulse dose of a solute (cobalt-EDTA) and a particle (chromium-mordanted fibre, <2 mm) marker with subsequent frequent faecal collection. Dry matter intake was 45.2 ± 8.0 g/kg^{0.75}/day. Mean retention times were 22.2 ± 5.3 h for solutes and 25.4 ± 5.2 h for particles, with the difference being not significant within individuals. This indicates the presence of a 'mucus-trap' colonic separation mechanism, which is in accord with morphological descriptions of the typical colonic furrow in chinchillas. Corresponding to a strategy of colonic digesta separation and caecotroph formation, secondary marker excretion peaks indicated coprophagic events that were spaced approximately 12 h apart. Given that these retention times appear longer than measures reported for rabbits (*Oryctolagus cuniculus*) or guinea pigs (*Cavia procillus*), it would be interesting to compare the digestive efficiency of chinchillas on high levels of dietary fibre to other species.

DOI: <https://doi.org/10.1111/jpn.12441>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-126021>

Journal Article

Accepted Version

Originally published at:

Hagen, K B; Dittmann, M T; Ortmann, S; Kreuzer, Michael; Hatt, Jean-Michel; Clauss, Marcus (2016). Retention of solute and particle markers in the digestive tract of chinchillas (*Chinchilla laniger*). *Journal of Animal Physiology and Animal Nutrition*, 100:801-806.

DOI: <https://doi.org/10.1111/jpn.12441>



Retention of solute and particle markers in the digestive tract of chinchillas (*Chinchilla laniger*)

Journal:	<i>Journal of Animal Physiology and Animal Nutrition</i>
Manuscript ID:	JAPAN-Jun-15-332.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Hagen, Katharina Dittmann, Marie; ETH Zürich, Department of Environmental Systems Science Ortmann, Sylvia Kreuzer, Michael; Institut für Nutztierwissenschaften, Tierernährung, ETH Zürich Hatt, Jean-Michel Clauss, Marcus; Division of Zoo Animals
Subject Area:	Exotics, Physiology, Rodents

SCHOLARONE™
Manuscripts

1 **Journal of Animal Physiology and Animal Nutrition - Revision**

2

3 **Retention of solute and particle markers in the digestive tract of**
4 **chinchillas (*Chinchilla laniger*)**

5

6 K. B. Hagen^{1,4}, M. T. Dittmann^{1,2}, S. Ortmann³, M. Kreuzer², J.-M. Hatt¹ and M. Clauss¹

7

8 ¹Clinic for Zoo Animals, Exotic Pets and Wildlife, Vetsuisse Faculty, University of Zurich,
9 Zurich, Switzerland

10 ²ETH Zurich, Institute for Agricultural Sciences, Zurich, Switzerland

11 ³Leibniz Institute for Zoo and Wildlife Research (IZW) Berlin, Germany

12 ⁴present address: Zoologischer Garten Halle GmbH, Halle (Saale), Germany

13

14 Correspondence: Marcus Clauss, Clinic for Zoo Animals, Exotic Pets and Wildlife, Vetsuisse
15 Faculty, University of Zurich, Winterthurerstr. 260, 8057 Zurich, Switzerland
16 mclauss@vetclinics.uzh.ch

17

18 Running head: Chinchilla digesta kinetics

19

Abstract

The chinchilla (*Chinchilla laniger*) is a herbivorous hystricomorph South American rodent for which no mean digesta retention times have been reported so far. Six animals (mean body mass \pm standard deviation: 513 ± 99 g) on a diet of grass hay and lucerne-based pellets were given a pulse dose of a solute (cobalt-EDTA) and a particle (chromium-mordanted fibre, <2 mm) marker with subsequent frequent faecal collection. Dry matter intake was 45.2 ± 8.0 g/kg^{0.75}/day. Mean retention times were 22.2 ± 5.3 h for solutes and 25.4 ± 5.2 h for particles, with the difference being not significant within individuals. This indicates the presence of a 'mucus trap' colonic separation mechanism, which is in accord with morphological descriptions of the typical colonic furrow in chinchillas. Corresponding to a strategy of colonic digesta separation and caecotroph formation, secondary marker excretion peaks indicated coprophagic events that were spaced approximately 12 h apart. Given that these retention times appear longer than measures reported for rabbits (*Oryctolagus cuniculus*) or guinea pigs (*Cavia procellus*), it would be interesting to compare the digestive efficiency of chinchillas on high levels of dietary fibre to other species.

Key words: chinchilla, digestion, coprophagy, caecotrophy, herbivory, fermentation

37 Introduction

38 Coprophagy is a digestive strategy employed by small herbivores to reduce potential losses of
39 nutrients associated with the shedding of microbial matter, originating from hindgut
40 fermentation, with the faeces (Karasov and Martínez del Río 2007). In order for this strategy
41 to be efficient, the material excreted as faeces has to be separated into the part that is foreseen
42 for re-ingestion, and the part that is definitely excreted. The mechanism that secures this
43 separation has been termed 'colonic separation mechanism' (CSM, Björnhag 1972). Two
44 fundamentally different CSM have been documented (Hume and Sakaguchi 1991; Björnhag
45 and Snipes 1999; Franz et al. 2011). The one present in lagomorphs (rabbits and hares) is
46 linked to a retrograde digesta washing in the colon that flushes bacteria back into the caecum,
47 and is called the 'wash back' CSM. The one present in rodents, called 'mucus trap' CSM, is
48 based on the extraction of microbes from the digesta into mucus contained in a specific
49 colonic structure, in which the trapped microbes are transported back to the caecum. In
50 hystricomorph ('porcupine-like', including guinea pigs *Cavia porcellus* or chinchillas)
51 rodents, this specific structure is the so-called colonic 'furrow' (Gorgas 1966; Snipes et al.
52 1988) and has been studied extensively (Takahashi and Sakaguchi 2000; 2006). In myomorph
53 ('mouse-like') rodents, the mechanism might be linked to certain longitudinal folds and
54 oblique furrows (*Plicae circulares*) in the colon (Behmann 1973; Sperber et al. 1983) but has
55 not been investigated in detail. From the caecum, the material separated by the CSM is
56 excreted as caecotrophs, re-ingested by the animal, and subject to enzymatic digestion in the
57 stomach and the small intestine. The corresponding mechanism in shrews, which also practice
58 coprophagy (Loxton et al. 1975), has not been identified.

59 Digesta passage experiments, in which an indigestible marker is ingested as a pulse-
60 dose by the animals, and its concentration subsequently documented in faeces over time, can
61 demonstrate coprophagic behaviour *per se* due to secondary marker peaks indicating repeated
62 marker uptake from the faeces (Clauss et al. 2007). Secondary peaks do not necessarily occur,

63 because reingestion of microbial components may not be required if the diet itself offers
64 nutrients, in particular protein, in high concentrations. Thus, coprophagy was shown to vary
65 with diet in rabbits (*Oryctolagus cuniculus*) (Fekete and Bokori 1985; García et al. 1995) or
66 viscachas (*Lagostomus maximus*) (Hagen et al. 2015a). Passage experiments can further
67 differentiate between the 'wash back' CSM, in which solute markers are particularly recycled
68 via coprophagy and hence have much longer retention times than particle markers, and the
69 'mucus trap' CSM in which solute and particle markers move together (Hume and Sakaguchi
70 1991; Pei et al. 2001; Franz et al. 2011).

71 Chinchillas (*Chinchilla laniger*) are herbivorous South American rodents. Their natural
72 diet mainly consists of grasses but also herbs, barks of bushes, bromeliads or fruits of
73 cactuses, depending on the availability of the respective plants (Cortés et al. 2002).
74 Chinchillas have a long history of being kept both as fur animals and as **pets**. **Feeding**
75 recommendations include the provision of a roughage source, high-fibre pellets and only
76 limited amounts of fresh vegetation or vegetables (Wolf et al. 2003; Grant 2014; Kohles
77 2014). Chinchillas have a colonic furrow typical for hystricomorph rodents (Gorgas 1966)
78 that is an integral part of their 'mucus-trap' CSM (Holtenius and Björnhag 1985). Without
79 indicating source or method, Johnson-Delaney (2006) states that the transit time of digesta
80 through the gastrointestinal tract of chinchillas is 12-15 h. An estimation for a mean digesta
81 **retention** time has been reported as 35 h in a study with food withdrawal, continuous marker
82 application for 24 h and long faecal sampling intervals (Krishnamurti et al. 1974), but so far,
83 the excretion patterns and mean retention times (MRT) for solute and particle markers have
84 not been investigated in this species. Such data is of special interest because, compared to
85 other hystricomorph rodents, chinchillas have a particularly long large intestine (Gorgas
86 1966). **This** may represent an adaptation to arid habitats and may allow them to live on lower
87 water intakes than other rodents (Hagen et al. 2014). We hypothesized that passage marker
88 excretion patterns would indicate coprophagy by secondary peaks, a 'mucus trap' CSM by

parallel movement of solute and particle markers, and yield long MRT values due to the long colon in this species.

91

92 **Material and Methods**

93 Six chinchillas (*Chinchilla laniger*; mean body mass \pm standard deviation: 513 \pm 99 g) were
94 used before in a study on food and water intake (Hagen et al., 2014) (experiment licence
95 80/2012). The animals were housed individually in open boxes with wood shavings in a room
96 with temperature ranging between 22 and 26 °C. The chinchillas were fed grass hay (g/kg dry
97 matter: crude protein, 221; neutral detergent fibre (NDF), 588) and commercial, lucerne-based
98 pellets (g/kg dry matter: crude protein, 139; NDF, 369) ad libitum. Food intake was quantified
99 by weighing the amount of diet items offered and weighing leftovers at the next feeding. The
100 MRT of a solute marker (cobalt-EDTA) and a particle marker (chromium-mordanted fibre
101 particles <2mm), prepared according to Udén et al. (1980), were measured. Chromium
102 concentration in the mordanted fibre was 37 g/kg dry matter. Markers were fed in quantities
103 of 0.01 g Co-EDTA and 0.04 g mordanted fibre per animal as a pulse-dose at 18:00 hrs,
104 mixed with a hay-based diet formula (Critical CareOxbow Animal Health, Murdock NE) and
105 administered orally with a syringe in manually restrained animals (taken out of their
106 enclosure). Care was taken to wipe off all non-ingested marker mix from the animals' snouts,
107 and to avoid carry-over of marker mix into the enclosure when placing the animals back.
108 Sampling of faeces was performed every 4 h during the first 48 h, every 6 h the next 48 h, and
109 once on day 5. Faeces were analysed for markers after wet ashing by atomic absorption
110 spectroscopy according to Behrend et al. (2004). Samples that deviated from the general
111 pattern of a decreasing marker concentration after an initial peak (i.e., samples later
112 interpreted as 'secondary peaks') were analysed in duplicate if sufficient amounts were
113 available. The MRT through the whole digestive tract were calculated according to
114 Thielemans et al. (1978) as

$$\text{MRT} = \frac{\sum t_i C_i dt_i}{\sum C_i dt_i}$$

with C_i = marker concentration in the fecal samples from the interval represented by time t_i (h after marker administration, using the midpoint of the sampling interval) and dt_i = the interval (h) of the respective sample

$$dt_i = \frac{(t_{i+1} - t_i) + (t_i - t_{i-1})}{2}$$

The marker was assumed to have been excreted completely once the fecal marker concentrations were similar to the background-levels determined in pre-dose fecal samples. The MRT for the two markers were normally distributed and compared by paired t-test using SPSS 21.0 (SPSS Inc. Chicago, IL).

Results

The relative average dry matter intake (per unit metabolic body weight) was 45.2 ± 8.1 (range 34-55) g/kg^{0.75}/day. Transit time (time of first marker appearance in faeces) was as short as 4 h in some individuals, and 8 or 12 h in the others (Fig. 1). The mean MRT for the solute marker (Co) was 22.2 ± 5.3 (range 16.3-30.2) h, and that of the particle marker (Cr) 25.4 ± 5.2 (range 19.5-34.1) h. The excretion patterns were characterised by 2 to 4 marker peaks (Fig. 1) that were sometimes only 12 h apart, in particular the first and the second peak. The two markers moved mostly in parallel. The difference between them was not significant at the 0.05 level even though a trend for a longer particle retention was apparent ($P = 0.082$). This was most likely due to two animals in which the majority of the particle marker was only excreted after the first coprophagic event (Fig. 1c and f).

Discussion

The results confirm that the chinchilla has a 'mucus-trap' CSM and practices coprophagy, a behaviour that is difficult to observe directly. With respect to the exact magnitude of digesta retention times in chinchillas, additional experiments using a variety of diets and intake levels are warranted. With respect to the markers used, it should be tested whether mordanted fibres are really representative for indigestible digesta in rodents, even if the widespread use of these markers facilitates the inclusion of the results of the present study in larger datasets (e.g. Müller et al. 2013). In particular, the identification of secondary marker peaks could be a bone of contention, because one could debate whether certain peaks are simply individual artefacts or truly represent coprophagy events. Care to avoid enclosure contamination with passage markers during marker application, and repeating analyses for critical samples, as both done in this study, can increase the interpretation safety only to a certain extent. Especially with respect to suggested secondary peaks with greater distance to the time of marker application, as e.g. marked in Fig. 1a, 1e or 1f, it would be beneficial to maintain the more frequent and labour-intensive faecal sampling regime that is typically only applied during the first two days in passage studies. Events that are not only indicated by a single peaking value, but by several samples indicating a recurrent increase and decrease in marker concentration (as e.g. evident in the first peaks in Fig. 1e), could more reliably be interpreted as non-artefacts. However, it must be noted that recurrent marker peaks have also been documented in individuals of species that are not known for coprophagy, where contamination of faeces with marker substrate could be excluded, and where no reingestion of faeces were observed during the experiment (e.g. Matsuda et al. 2015). Therefore, marker excretion patterns should always be considered in the context of other morphological and physiological data.

Given the variation in whether similar or dissimilar concentrations of both markers were excreted at secondary peaks, it must be concluded that the 'mucus-trap' CSM shows no clear preference for either the solute or the particle marker, and that possibly a degree of chance is

involved with respect to the degree a marker is retained. The spacing of the repeated marker excretion peaks suggests a rhythm of coprophagy that is shorter than 24 h and closer to 12 h. This matches the findings of Holtenius and Björnhag (1985) who observed single coprophagy events in chinchillas during the whole day (but not during the night). Similarly, the distance between marker excretion peaks in viscachas suggested coprophagy events at 12 h time intervals (Clauss et al. 2007). Also, some distances between marker excretion peaks in rabbits and guinea pigs were of similar magnitude (Sakaguchi et al. 1987; Franz et al. 2011). However, in rabbits, marker peaks spaced 24 h apart have also been reported (Sakaguchi et al. 1987). Apparently, the rhythm of coprophagic events varies between species. Some rodents practice coprophagy multiple times during the day, such as the herbivorous vole (*Microtus californicus*) (Kenagy and Hoyt 1979), the meadow vole (*M. pennsylvanicus*) (Ouellette and Heisinger 1980), the Norway lemming (*Lemmus lemmus*) (Björnhag 1977) or the mountain beaver (*Aplodontia rufa*) (Ingles 1961). The degu (*Octodon degu*) is another animal that practices coprophagy about every 12 h, but in contrast to the chinchilla, the degu shows this behaviour only at night (Kenagy et al. 1999). In mountain hares (*Lepus timidus*) (Pehrson 1983), kangaroo rats (*Dipodomys microps*) (Kenagy and Hoyt 1979), nutrias (*Myocastor coypus*) (Gosling 1979) and capybaras (*Hydrochoerus hydrochaeris*) (Herrera 1985), a rhythm of 24 h for coprophagy was observed. Apart from these apparent species differences, variation in the nutrient composition and the amount of the available diet may change coprophagic behaviour within species (Fekete and Bokori 1985; García et al. 1995; Kenagy et al. 1999; Hagen et al. 2015a).

In contrast to MRTs in rabbits (app. 15 h) and guinea pigs (app. 18 h) with the same passage markers on a grass hay-only diet (Franz et al. 2011), chinchillas appear to achieve surprisingly long MRTs for their body size, reaching values in the lower range of horses on a grass hay-only diet with the same marker system (23-34 h, Clauss et al. 2014). Whether this translates into higher fibre digestibilities in chinchillas than for example in rabbits and guinea

pigs or is simply a side effect of the fact that chinchillas have, for hystricomorph rodents, a particularly long large intestine for water re-absorption as an adaptation to their arid habitat (Gorgas, 1966), remains to be investigated. The few digestibility measurements that have been reported for chinchillas do not allow to conclusively decide whether the digestive efficiency of chinchillas is equally or less affected by dietary fibre compared to rabbits or guinea pigs (Hagen et al. 2015b). Reports that pet chinchillas suffer from body mass losses when kept on a diet of either grass hay or fresh grass alone (Wolf et al. 2003) apparently do not match a high fibre digestibility one would intuitively link with long retention times. Even though coprophagy is understood mainly as a mechanism to recycle nutrients related to gastrointestinal microbes (reviewed in Karasov and Martínez del Río 2007; Franz et al. 2011), the fact that the fibre marker in the present study and similar studies is also recycled by this mechanism suggests that some fibrous components will also be submitted to a second, and hence prolonged, fermentation in the gastrointestinal tract of hindgut fermenters with a 'mucus trap' CSM.

Acknowledgements

We thank Professor Max M. Wittenbrink for the use of the holding facilities, Sina Bigger for lending us the chinchillas, Heidrun Barleben and Carmen Kunz for sample analysis, and two anonymous reviewers for comments on the manuscript. This study was part of project 310030_135252/1 funded by the Swiss National Science Foundation.

Literature

Behmann, H., 1973: Vergleichend- und funktionell-anatomische Untersuchungen am Caecum und Colon myomorpher Nagetiere. *Zeitschrift für wissenschaftliche Zoologie* **186**, 173-294.

- 214 Behrend, A.; Lechner-Doll, M.; Streich, W.J.; Clauss, M., 2004: Seasonal faecal excretion,
215 gut fill, liquid and particle marker retention in mouflon (*Ovis ammon musimon*), and a
216 comparison with roe deer (*Capreolus capreolus*). *Acta Theriologica* **49**, 503-515.
- 217 Björnhag, G., 1977: Demonstration of coprophagy in some rodents. *Swedish Journal of*
218 *Agricultural Research* **7**, 105-113.
- 219 Björnhag, G., 1972: Separation and delay of contents in the rabbit colon. *Swedish Journal of*
220 *Agricultural Research* **2**, 125-136.
- 221 Björnhag, G.; Snipes, R.L., 1999: Colonic separation mechanism in lagomorph and rodent
222 species - a comparison. *Zoosystematics and Evolution* **75**, 275-281.
- 223 Clauss, M.; Besselmann, D.; Schwarm, A.; Ortmann, S.; Hatt, J.-M., 2007: Demonstrating
224 coprophagy with passage markers? The example of the plains viscacha (*Lagostomus*
225 *maximus*). *Comparative Biochemistry and Physiology A* **147**, 453-459.
- 226 Clauss, M.; Schiele, K.; Ortmann, S.; Fritz, J.; Codron, D.; Hummel, J.; Kienzle, E., 2014:
227 The effect of very low food intake on digestive physiology and forage digestibility in horses.
228 *Journal of Animal Physiology and Animal Nutrition* **98**, 107-118.
- 229 Cortés, A.; Miranda, E.; Jiménez, J.E., 2002: Seasonal food habits of the endangered long-
230 tailed chinchilla (*Chinchilla lanigera*): the effect of precipitation. *Mammalian Biology* **67**,
231 167-175.
- 232 Fekete, S.; Bokori, J., 1985: The effect of the fiber and protein level of the ration upon the
233 cecotrophy of rabbit. *Journal of Applied Rabbit Research* **8**, 68-71.
- 234 Franz, R.; Kreuzer, M.; Hummel, J.; Hatt, J.-M.; Clauss, M., 2011: Intake, selection, digesta
235 retention, digestion and gut fill of two coprophageous species, rabbits (*Oryctolagus*
236 *cuniculus*) and guinea pigs (*Cavia porcellus*), on a hay-only diet. *Journal of Animal*
237 *Physiology and Animal Nutrition* **95**, 564-570.
- 238 García, J.; De Blas, J.C.; Carabaño, R.; Garcia, P., 1995: Effect of type of lucerne hay on
239 caecal fermentation and nitrogen contribution through caecotrophy in rabbits. *Reproduction*
240 *Nutrition Development* **35**, 267-275.

- 241 Gorgas, M., 1966: Vergleichend-anatomische Untersuchungen am Magen-Darm-Kanal der
242 Sciuromorpha, Hystricomorpha und Caviomorpha (Rodentia). *Zeitschrift für*
243 *wissenschaftliche Zoologie* **175**, 237-404.
- 244 Gosling, L.M., 1979: The twenty-four hour activity cycle of captive coypus (*Myocastor*
245 *coypus*). *Journal of Zoology* **187**, 341-367.
- 246 Grant, K., 2014: Rodent nutrition: digestive comparisons of 4 common rodent species.
247 *Veterinary Clinics of North America: Exotic Animal Practice* **17**, 471-483.
- 248 Hagen, K.; Clauss, M.; Hatt, J.-M., 2014: Drinking preferences in chinchillas (*Chinchilla*
249 *laniger*), degus (*Octodon degu*) and Guinea pigs (*Cavia porcellus*). *Journal of Animal*
250 *Physiology and Animal Nutrition* **98**, 942-947.
- 251 Hagen, K.B.; Besselmann, D.; Cyrus-Eulenberger, U.; Vendl, C.; Ortmann, S.; Zingg, R.;
252 Kienzle, E.; Kreuzer, M.; Hatt, J.-M.; Clauss, M., 2015a: Digestive physiology of the plains
253 viscacha (*Lagostomus maximus*), a large herbivorous hystricomorph rodent. *Zoo Biology*
254 **(online) doi 10.1002/zoo.21216**.
- 255 Hagen, K.B.; Tschudin, A.; Liesegang, A.; Hatt, J.-M.; Clauss, M., 2015b: Organic matter
256 and macromineral digestibility in domestic rabbits (*Oryctolagus cuniculus*) as compared to
257 other hindgut fermenters. *Journal of Animal Physiology and Animal Nutrition (online) doi*
258 **10.1111/jpn.12323**.
- 259 Herrera, E.A., 1985: Coprophagy in the capybara, *Hydrochoerus hydrochoeris*. *Journal of*
260 *Zoology* **207**, 616-619.
- 261 Holtenius, K.; Björnhag, G., 1985: The colonic separation mechanism in the guinea pig
262 (*Cavia porcellus*) and the chinchilla (*Chinchilla laniger*). *Comparative Biochemistry and*
263 *Physiology A* **82**, 537-542.
- 264 Hume, I.D.; Sakaguchi, E., 1991: Patterns of digesta flow and digestion in foregut and
265 hindgut fermenters, In: Tsuda, T.; Saaski, Y.; Kawashima, R. (eds.), *Physiological aspects of*
266 *digestion and metabolism in ruminants*. Academic Press, San Diego. 427-451.
- 267 Ingles, L.G., 1961: Reingestion in the mountain beaver. *Journal of Mammalogy* **42**, 411-412.
- 268 Johnson-Delaney, C.A., 2006: Anatomy and physiology of the rabbit and rodent
269 gastrointestinal system. *Proceedings of the Association of Avian Veterinarians* **2006**, 9-17.

- 270 Karasov, W.H.; Martínez del Rio, C., 2007: *Physiological ecology: how animals process*
271 *energy, nutrients, and toxins*. Princeton University Press, Princeton, New Jersey, USA.
- 272 Kenagy, G.J.; Hoyt, D., 1979: Reingestion of feces in rodents and its daily rhythmicity.
273 *Oecologia* **44**, 403-409.
- 274 Kenagy, G.J.; Veloso, C.; Bozinovic, F., 1999: Daily rhythms of food intake and feces
275 reingestion in the degu, an herbivorous chilean rodent: optimizing digestion through
276 coprophagy. *Physiological and Biochemical Zoology* **72**, 78-86.
- 277 Kohles, M., 2014: Gastrointestinal anatomy and physiology of select exotic companion
278 mammals. *Veterinary Clinics of North America: Exotic Animal Practice* **17**, 165-178.
- 279 Krishnamurti, C.R.; Kitts, W.D.; Smith, D.C., 1974: The digestion of carbohydrates in the
280 chinchilla (*Chinchilla lanigera*). *Canadian Journal of Zoology* **52**, 1227-1233.
- 281 Loxton, R.G.; Raffaelli, D.; Begon, M., 1975: Coprophagy and the diurnal cycle of the
282 common shrew, *Sorex araneus*. *Journal of Zoology* **177**, 449-453.
- 283 Matsuda, I.; Sha, J.C.M.; Ortmann, S.; Schwarm, A.; Grandl, F.; Caton, J.; Jens, W.; Kreuzer,
284 M.; Marlena, D.; Hagen, K.B.; Clauss, M., 2015: Excretion patterns of solute and different-
285 sized particle passage markers in foregut-fermenting proboscis monkey (*Nasalis larvatus*) do
286 not indicate an adaptation for rumination. *Physiology and Behavior* **149**, 45-52.
- 287 Müller, D.W.H.; Codron, D.; Meloro, C.; Munn, A.; Schwarm, A.; Hummel, J.; Clauss, M.,
288 2013: Assessing the Jarman-Bell Principle: scaling of intake, digestibility, retention time and
289 gut fill with body mass in mammalian herbivores. *Comparative Biochemistry and Physiology*
290 *A* **164**, 129-140.
- 291 Ouellette, D.E.; Heisinger, J.F., 1980: Reingestion of feces by *Microtus pennsylvanicus*.
292 *Journal of Mammalogy* **61**, 366-368.
- 293 Pehrson, Å., 1983: Caecotrophy in caged mountain hares (*Lepus timidus*). *Journal of Zoology*
294 **199**, 563-574.
- 295 Pei, Y.X.; Wang, D.H.; Hume, I.D., 2001: Selective digesta retention and coprophagy in
296 Brandt's vole (*Microtus brandti*). *Journal of Comparative Physiology B* **171**, 457-464.

- 297 Sakaguchi, E.; Itoh, H.; Uchida, S.; Horigome, T., 1987: Comparison of fibre digestion and
298 digesta retention time between rabbits, guinea-pigs, rats and hamsters. *British Journal of*
299 *Nutrition* **58**, 149-158.
- 300 Snipes, R.L.; Hörnicke, H.; Björnhag, G.; Stahl, W., 1988: Regional differences in hindgut
301 structure and function in the nutria (*Myocastor coypus*). *Cell Tissue Research* **252**, 435-447.
- 302 Sperber, I.; Björnhag, G.; Ridderstråle, Y., 1983: Function of proximal colon in lemming and
303 rat. *Swedish Journal of Agricultural Research* **13**, 243-256.
- 304 Takahashi, T.; Sakaguchi, E., 2000: Role of the furrow of the proximal colon in the
305 production of soft and hard feces in nutrias (*Myocastor coypus*). *Journal of Comparative*
306 *Physiology B* **170**, 531-535.
- 307 Takahashi, T.; Sakaguchi, E., 2006: Transport of bacteria across and along the large intestinal
308 lumen of guinea pigs. *Journal of Comparative Physiology B* **176**, 173-178.
- 309 Thielemans, M.F.; Francois, E.; Bodart, C.; Thewis, A., 1978: Mesure du transit
310 gastrointestinal chez le porc a l'aide des radiolanthanides. Comparaison avec le mouton.
311 *Annales De Biologie Animale, Biochimie, Biophysique* **18**, 237-247.
- 312 Wolf, P.; Schröder, A.; Wenger, A.; Kamphues, J., 2003: The nutrition of the chinchilla as a
313 companion animal - basic data, influences and dependences. *Journal of Animal Physiology*
314 *and Animal Nutrition* **87**, 129-133.

315

316 **Figure 1** Marker excretion patterns for a solute (cobalt-EDTA) and a particle marker
317 (chromium-mordanted fibre, < 2 mm) in the six chinchillas (*Chinchilla laniger*, a-e) of this
318 study. Note the recurrent marker peaks (marked by arrows), indicative of coprophagy, and the
319 general absence of difference in pattern between the two markers within individuals.
320

For Peer Review




Figure 1 Marker excretion patterns for a solute (cobalt-EDTA) and a particle marker (chromium-mordanted fibre, < 2 mm) in the six chinchillas (*Chinchilla laniger*) of this study. Note the recurrent marker peaks (marked by arrows), indicative of coprophagy, and the general absence of difference in pattern between the two markers within individuals.




Figure 1 Marker excretion patterns for a solute (cobalt-EDTA) and a particle marker (chromium-mordanted fibre, < 2 mm) in the six chinchillas (*Chinchilla laniger*, a-e) of this study. Note the recurrent marker peaks (marked by arrows), indicative of coprophagy, and the general absence of difference in pattern between the two markers within individuals.

13411x10617mm (1 x 1 DPI)




Figure 1 Marker excretion patterns for a solute (cobalt-EDTA) and a particle marker (chromium-mordanted fibre, < 2 mm) in the six chinchillas (*Chinchilla laniger*, a-e) of this study. Note the recurrent marker peaks (marked by arrows), indicative of coprophagy, and the general absence of difference in pattern between the two markers within individuals.

13385x10591mm (1 x 1 DPI)




Figure 1 Marker excretion patterns for a solute (cobalt-EDTA) and a particle marker (chromium-mordanted fibre, < 2 mm) in the six chinchillas (*Chinchilla laniger*, a-e) of this study. Note the recurrent marker peaks (marked by arrows), indicative of coprophagy, and the general absence of difference in pattern between the two markers within individuals.

13436x10668mm (1 x 1 DPI)




Figure 1 Marker excretion patterns for a solute (cobalt-EDTA) and a particle marker (chromium-mordanted fibre, < 2 mm) in the six chinchillas (*Chinchilla laniger*, a-e) of this study. Note the recurrent marker peaks (marked by arrows), indicative of coprophagy, and the general absence of difference in pattern between the two markers within individuals.




Figure 1 Marker excretion patterns for a solute (cobalt-EDTA) and a particle marker (chromium-mordanted fibre, < 2 mm) in the six chinchillas (*Chinchilla laniger*, a-e) of this study. Note the recurrent marker peaks (marked by arrows), indicative of coprophagy, and the general absence of difference in pattern between the two markers within individuals.

13411x10617mm (1 x 1 DPI)




Figure 1 Marker excretion patterns for a solute (cobalt-EDTA) and a particle marker (chromium-mordanted fibre, < 2 mm) in the six chinchillas (*Chinchilla laniger*, a-e) of this study. Note the recurrent marker peaks (marked by arrows), indicative of coprophagy, and the general absence of difference in pattern between the two markers within individuals.